

ESTD 1962

Kendamil®

healthcare

Classic Toddler Milk 800g

Kendamil Classic Toddler Milk is a formula uniquely made with whole milk and free from palm oil, fish oil, and soy, formulated to help meet the nutritional needs of toddlers aged 1-3 years as part of a varied, balanced diet. Kendamil is enriched with plant-based DHA and ARA, making it the only vegetarian friendly baby milk. Kendamil Toddler Milk contains a blend of GOS and HMOs in the form of 3'-GL (galactosylactose).



Whole milk from
grass-fed cows



Plant-based DHA
(Omega-3)



No palm oil.
No fish oil.
Vegetarian friendly

Presentation

Kendamil Stage 3 Classic is packaged in a 800g can.

Contraindications

Not suitable for infants with galactosaemia, lactose intolerance, or allergy to any of the ingredients.

Shelf life

24 months from date of manufacture.

Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

IMPORTANT NOTICE

This information is intended for Healthcare Professional use only. Toddler milk should be used as part of a varied, balanced diet from 1 year.

Ingredients

Whole **Milk**, Skimmed **Milk**, Lactose (from **Milk**), Vegetable Oils (Sunflower, Coconut, Rapeseed), Galacto-oligosaccharides (**Milk**) Demineralised Whey (from **Milk**), **Less than 2% of:** Calcium Phosphate, Potassium Hydroxide, Oil from the Microalgae Schizochytrium sp., Oil from Mortierella Alpina, Sodium Ascorbate, Fructo-oligosaccharides, Calcium Citrate, Iron Pyrophosphate, Nucleotides (cytidine-5'-monophosphate, disodium uridine-5'-monophosphate, adenosine-5'-monophosphate, disodium inosine-5'-monophosphate, disodium guanosine-5'-monophosphate), Vitamin E, Zinc Sulphate, Niacin, Calcium Pantothenate, Copper Sulphate, Vitamin A, Thiamin, Riboflavin, Vitamin B6, Folic Acid, Manganese Sulphate, Potassium Iodide, Vitamin K, Sodium Selenite, Biotin, Vitamin D3, Vitamin B12, Antioxidant: Tocopherol-Rich Extract.

Allergy Advice: For Allergens See Ingredients in **Bold**.

Nutritional Values

		Per 100g	Per 100ml	Vitamin C	mg	86	12 (27%*)
Energy	kJ kcal	1974 471	270 65	Thiamin	mg	0.52	0.07 (14%*)
Fat	g	20.4	2.8	Riboflavin	mg	1.1	0.15 (21%*)
of which saturates	g	9.3	1.3	Niacin	mg	5.2	0.71 (10%*)
of which unsaturates	g	11.1	1.5	Vitamin B6	mg	0.32	0.04
Omega-3				Folate (DFE)	mcg	100	14 (19%*)
α-Linolenic Acid (ALA)	mg	260	36	Vitamin B12	mcg	1.3	0.18 (23%*)
Docosahexaenoic Acid (DHA)	mg	62	8.5	Pantothenic Acid	mg	3.2	0.44 (15%*)
Omega 6				Biotin	mcg	13	1.8 (18%*)
Linoleic Acid (LA)	mg	2600	356	Minerals			
Arachidonic Acid (ARA)	mg	16	2.2	Sodium	mg	170	23
Total Carbohydrate	g	58	7.9	Potassium	mg	610	84 (8%*)
of which sugars*	g	54	7.4	Chloride	mg	392	54 (11%*)
of which lactose	g	53	7.3	Calcium	mg	880	121 (22%*)
Fibre	g	1.66	0.23	Phosphorous	mg	480	66 (12%*)
of which Galacto-oligosaccharides	g	1.45	0.2	Magnesium	mg	47	6.4 (8%*)
of which 3'+4'+6'-GL**	g	0.18	0.02	Iron	mg	6.3	0.9 (11%*)
of which Fructo-oligosaccharides	g	0.16	0.02	Zinc	mg	3.7	0.5 (10%*)
Protein (N x 6.25)	g	13	1.8	Copper	mcg	0.26	0.04 (8%*)
Salt	g	0.42	0.06	Iodine	mcg	115	16 (20%*)
Vitamins				Selenium	mcg	18	2.5 (13%*)
Vitamin A	mcg-RE	450	62 (16%*)	Manganese	mcg	65	8.9
Vitamin D3	mcg	5.4	0.7 (10%*)	Others			
Vitamin E	mg-α-TE	9	1.2 (24%*)	L-Carnitine (naturally occurring)	mg	6.8	0.9
Vitamin K	mcg	27	3.7 (31%*)	Nucleotides	mg	20	2.7

*Sugars derived from milk **GL - Galactosyllactose

Feeding Guide (12-36 months)

(1 level scoop = 4.6g) Add 1 level scoop of Kendamil 3 powder to each 30ml of water.

Amount of freshly boiled and cooled water per feed	Number of scoops per feed
180ml	6

Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist. If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.