



Goat Follow-On Milk 800g

Kendamil Goat Follow On Milk is a formula uniquely made with whole goat milk and free from palm oil, fish oil and soy, formulated to help meet the increased nutritional needs of infants from 6 months of age alongside a weaning diet. Kendamil is enriched with plant-based DHA and ARA, making it vegetarian friendly. Kendamil Follow On Milk contains a blend of GOS and HMOs in the form of 3'-GL (galactosyllactose)



Made with creamy
Whole A2 Goat Milk
and Goat Whey



No palm oil.
No fish oil.
Vegetarian friendly



Contains GOS
and free from
maltodextrin

Presentation

Kendamil Stage 2 Goat is packaged in a 800g can.

Contraindications

Not suitable for those with allergy to any of the ingredients, including cows milk proteins.

Shelf life

24 months from date of manufacture.

Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

IMPORTANT NOTICE

Important notice: This information is intended for Healthcare Professional use only. Breastfeeding is best for babies. Follow-on milk is only for babies over 6 months, as part of a mixed diet and should not be used as a breastmilk substitute before 6 months. The use of follow-on formula should be made only on the advice of a healthcare professional.

Ingredients

Whole Goat **Milk**, Demineralised Goat Whey Powder (from **Milk**), Vegetable Oils (Sunflower, Coconut, Rapeseed), Lactose (from **Milk**), Galacto-oligosaccharides (from **Milk**), **Less than 2% of:** TriCalcium Citrate, Choline Bitartrate, Sodium Citrate, Potassium Chloride, Oil from the microalgae Schizochytrium sp. (**Milk**), L-Phenylalanine, Magnesium Chloride, L-Isoleucine, L-Tryptophan, Potassium Hydroxide, Ascorbic acid, Iron Pyrophosphate, Oil from Mortierella Alpina (**Milk**), L-Tyrosine, Inositol, Taurine, TriCalcium Phosphate, Calcium Phosphate, Nucleotides (Cytidine-5'-Monophosphate, Disodium Uridine-5'-Monophosphate, Adenosine-5'-Monophosphate, Disodium Inosine-5'-Monophosphate, Disodium Guanosine-5'-Monophosphate), Zinc Sulphate, Vitamin E, Niacin, Calcium Pantothenate, Copper Sulphate, Thiamin, Riboflavin, Vitamin A, Vitamin B6, Manganese Sulphate, Folic Acid, Potassium Iodide, Sodium Selenite, Vitamin K, Vitamin D3, Biotin, Vitamin B12.

Allergy Advice: For Allergens See Ingredients In **Bold**.

Nutritional Values

		Per 100g	Per 100ml	Vitamin B6	mg	0.41	0.05
Energy	kJ kcal	2137 511	276 66	Folate (DFE)	mcg	133	17
Fat	g	26.8	3.5	Vitamin B12	mcg	0.9	0.12
of which saturates	g	9.8	1.2	Pantothenic Acid	mg	3.7	0.5
of which unsaturates	g	17	2.2	Biotin	mcg	24	3.1
Omega-3				Minerals			
α-Linolenic Acid (ALA)	mg	365	47	Sodium	mg	170	22
Docosahexaenoic Acid (DHA - LCP)	mg	120	15	Potassium	mg	660	85
Omega 6				Chloride	mg	500	64
Linoleic Acid (LA)	mg	3650	471	Calcium	mg	340	44
Arachidonic Acid (AA - LCP)	mg	60	7.7	Phosphorous	mg	200	26
Total Carbohydrate	g	56	7.2	Magnesium	mg	48	6.2
of which sugars*	g	54	7	Iron	mg	6.3	0.81
of which lactose	g	52	6.7	Zinc	mg	4.3	0.55
Fibre	g	1.4	0.18	Copper	mg	0.45	0.058
of which Galacto-oligosaccharides	g	1.35	0.17	Iodine	mcg	110	14
of which 3'GL**	g	0.05	0.01	Selenium	mcg	20	2.6
Protein (N x 6.25)	g	10.5	1.4	Manganese	mcg	60	7.7
Vitamins				Fluoride	mg	<0.1	<0.015
Vitamin A	mcg-RE	441	57	Others			
Vitamin D3	mcg	11	1.4	Taurine	mg	45	5.8
Vitamin E	mg-α-TE	17	2.2	Choline	mg	125	16
Vitamin K	mcg	32	4.1	Inositol	mg	80	10
Vitamin C	mg	92	11.9	L-Carnitine	mg	9	1
Thiamin	mg	0.5	0.06	Nucleotides	mg	22	2.8
Riboflavin	mg	1.2	0.15				
Niacin	mg	5.2	0.7				

*Sugars derived from milk **3'GL - Galactosyllactose

Feeding Guide (6-12 months)

(1 level scoop = 4.3g) Add 1 level scoop of powder to each 30ml of water^a. This feeding table should only be used as a guide. All babies are different and so may require more or less than is shown below. Kendamil Goat Follow-on Milk should be fed on demand. If you need advice talk to your healthcare professional.

Water per feed ^a	Scoops per feed ^b
150ml	5
180ml	6
210ml	7

Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist. If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.