

ESTD 1962

# Kendamil®

healthcare

## Goat Toddler Milk 800g

Kendamil Goat Toddler Milk is a formula uniquely made with whole goat milk and free from palm oil, fish oil and soy, formulated to help meet the nutritional needs of toddlers aged 1-3 years as part of a varied, balanced diet. Kendamil is enriched with plant-based DHA and ARA, making it vegetarian friendly. Kendamil Toddler Milk contains a blend of GOS and HMOs in the form of 3'-GL (galactosyllactose)



Made with creamy  
Whole A2 Goat Milk  
and Goat Whey



No palm oil.  
No fish oil.  
Vegetarian friendly



Contains GOS  
and free from  
maltodextrin

### Presentation

Kendamil Stage 3 Goat is packaged in a 800g can.

### Contraindications

Not suitable for those with allergy to any of the ingredients, including cows milk proteins.

### Shelf life

24 months from date of manufacture.

### Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

### IMPORTANT NOTICE

Important notice: This information is intended for Healthcare Professional use only. Toddler milk should be used as part of a varied, balanced diet from 1 year.

### Ingredients

Whole goat **milk**, Goat demineralised whey powder (from **milk**), Lactose (from **milk**), Vegetable oils (sunflower, coconut, rapeseed), Galacto-oligosaccharides (from **milk**), TriCalcium citrate, TriCalcium Phosphate, Calcium phosphate, Potassium hydroxide, Ascorbic acid, Iron Pyrophosphate, Oil from the microalgae Schizochytrium sp. (**milk**), Oil from Mortierella alpina (**milk**), Nucleotides (Cytidine-5'-monophosphate, disodium Uridine-5'-monophosphate, Adenosine-5'-monophosphate, Disodium Inosine-5'-monophosphate, Disodium Guanosine-5'-monophosphate), Vitamin E, Zinc sulphate, Niacin, Calcium pantothenate, Copper sulphate, Vitamin A, Thiamin, Riboflavin, Vitamin B6, Folic acid, Manganese sulphate, Potassium iodide, Sodium selenite, Vitamin K, Biotin, Vitamin D3, Vitamin B12. **Allergy Advice:** For Allergens See Ingredients In **Bold**.

## Nutritional Values

		Per 100g	Per 100ml				
				Vitamin C	mg	95	13 (29%*)
Energy	kJ kcal	1980 473	273 65	Thiamin	mg	0.55	0.08 (16%*)
Fat	g	21.3	2.9	Riboflavin	mg	0.8	0.11 (15%*)
of which saturates	g	10.8	1.5	Niacin	mg	5.8	0.80 (11%*)
of which unsaturates	g	10.5	1.4	Vitamin B6	mg	0.36	0.05
<b>Omega-3</b>				Folate (DFE)	mcg	109	15 (20%*)
α-Linolenic Acid (ALA)	mg	190	26	Vitamin B12	mcg	0.8	0.11 (14%*)
Docosahexaenoic Acid (DHA - LCP)	mg	68	9.4	Pantothenic Acid	mg	3.6	0.5 (17%*)
<b>Omega 6</b>				Biotin	mcg	14	2.0 (20%*)
Linoleic Acid (LA)	mg	1800	248	<b>Minerals</b>			
Arachidonic Acid (AA - LCP)	mg	36	5	Sodium	mg	180	25
Total Carbohydrate	g	57	7.9	Potassium	mg	900	124 (13%*)
of which sugars*	g	55	7.6	Chloride	mg	600	83 (16%*)
of which lactose	g	53	7.3	Calcium	mg	780	108 (20%*)
Fibre	g	1.5	0.21	Phosphorous	mg	500	69 (12%*)
of which Galacto-oligosaccharides	g	1.45	0.2	Magnesium	mg	50	6.9 (8%*)
of which 3'GL**	g	0.05	0.01	Iron	mg	7	1 (13%*)
Protein (N x 6.25)	g	13	1.8	Zinc	mg	4	0.5 (10%*)
Salt	g	0.46	0.06	Copper	mcg	0.28	0.040 (8%*)
<b>Vitamins</b>				Iodine	mcg	90	13 (16%*)
Vitamin A	mcg-RE	500	69 (17%*)	Selenium	mcg	20	3 (15%*)
Vitamin D3	mcg	6	0.8 (11%*)	Manganese	mcg	67	9.2
Vitamin E	mg-α-TE	10	1.4 (28%*)	<b>Others</b>			
Vitamin K	mcg	30	4.1 (34%*)	L-Carnitine	mg	9	1.3
				Nucleotides	mg	20	2.8

\*Sugars derived from milk \*\*3'GL - Galactosyllactose

### Feeding Guide (12-36 months)

(1 level scoop = 4.6g) Add 1 level scoop of powder to each 30 ml of water.

Amount of freshly boiled and cooled water per feed	Number of scoops per feed
180ml	6

### Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist. If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.

For healthcare professional use.