

ESTD 1962

Kendamil®

healthcare

Organic Toddler Milk 800g

Kendamil Organic Toddler Milk is a formula uniquely made with organic whole milk and free from palm oil, fish oil and soy. It's formulated to help meet the nutritional needs of toddlers aged 1-3 years as part of a varied, balanced diet. Exclusively manufactured in the UK and certified organic by the Soil Association,

Kendamil is enriched with plant-based DHA and ARA, making it the only vegetarian friendly baby milk. Kendamil Toddler Milk contains a blend of GOS and HMOs in the form of 3'-GL (galactosylactose).



Made in the UK



Organic whole milk from grass-fed British family farms



No palm oil.
No fish oil.
Vegetarian friendly

Presentation

Kendamil Stage 3 Organic is packaged in a 800g can.

Contraindications

Not suitable for those with allergies to any of the ingredients, galactosaemia or cow's milk intolerance.

Shelf life

24 months from date of manufacture.

Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

IMPORTANT NOTICE

This information is intended for Healthcare Professional use only. Toddler milk should be used as part of a varied, balanced diet from 1 year.

Ingredients

Organic whole **milk**, Organic lactose (from **milk**), Organic whole **milk** powder, Organic vegetable oils (sunflower, coconut, rapeseed), Organic skimmed **milk** powder, Organic demineralised **milk** whey protein powder, Organic galacto-oligosaccharides (from **milk**), **Less than 2% of:** Calcium lactate, Oil from the microalgae Schizochytrium sp., Calcium phosphate, Potassium hydroxide, Sodium Citrate, Vitamin C, Calcium citrate, Potassium chloride, Iron pyrophosphate, Vitamin E, Zinc sulphate, Niacin, Calcium Pantothenate, Copper sulphate, Thiamin, Riboflavin, Vitamin A, Vitamin B6, Manganese sulphate, Folic acid, Sodium selenite, Potassium Iodide, Vitamin K, Vitamin D3, Biotin, Vitamin B12.

For Allergens: See ingredients in **Bold**

Nutritional Values

		Per 100g	Per 100ml				
				Riboflavin	mg	0.8	0.11
Energy	kJ kcal	2033 485	279 67	Niacin	mg	4.7	0.64
Fat	g	23.2	3.2	Vitamin B6	mg	0.4	0.05
of which saturates	g	9.3	1.3	Folate (DFE)	mcg	150	20.6
of which unsaturates	g	12.6	1.7	Vitamin B12	mcg	1.5	0.21
Omega-3				Pantothenic Acid	mg	3.8	0.52
α-Linolenic Acid (ALA)	mg	320	44	Biotin	mcg	12	1.64
Docosahexaenoic Acid (DHA)	mg	100	13.7	Minerals			
Omega 6				Sodium	mg	170	23
Linoleic Acid (LA)	mg	3250	445	Potassium	mg	700	96
Total Carbohydrate	g	57	7.8	Chloride	mg	320	44
of which sugars*	g	55.7	7.6	Calcium	mg	540	74
of which lactose	g	53.9	7.4	Phosphorous	mg	320	44
Fibre	g	1.5	0.21	Magnesium	mg	25	3.4
of which Galacto-oligosaccharides	g	1.45	0.2	Iron	mg	6.2	0.85
of which 3'+4'+6'-GL**	g	0.18	0.02	Zinc	mg	3.8	0.52
Protein (N x 6.25)	g	11.4	1.6	Copper	mg	0.32	0.04
Vitamins				Iodine	mcg	90	12.3
Vitamin A	mcg-RE	380	52	Selenium	mcg	15	2.1
Vitamin D3	mcg	10	1.4	Manganese	mcg	65	8.9
Vitamin E	mg-α-TE	9	1.2	Fluoride	mg	<0.1	<0.015
Vitamin K	mcg	25	3.4	Others			
Vitamin C	mg	70	9.6	L-Carnitine (naturally present)	mg	12	1.6
Thiamin	mg	0.4	0.05				

*Sugars derived from milk **GL - Galactosyllactose

Feeding Guide (12-36 months)

(1 level scoop = 4.6g) Add 1 level scoop of Kendamil Organic 3 powder to each 30ml of water.

Amount of freshly boiled and cooled water per feed	Number of scoops per feed
180ml	6

Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist. If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.