



ESTD 1942

for healthcare professional use

Kendamil®

product data card

Classic Follow-On Milk 800g

Description

Kendamil Classic Follow-On Milk is formula uniquely made with whole milk fat nutrients and free from palm oil and fish oil, formulated to help meet the increased nutritional needs of infants from 6 months of age alongside a weaning diet. Exclusively manufactured in the UK using Red-Tractor certified whole milk, Kendamil is enriched with plant-based DHA and ARA, making it the only vegetarian friendly baby milk. Kendamil Follow-On Milk contains a blend of GOS/FOS and HMOs in the form of 3'-GL (galactosyllactose).

Presentation

Kendamil Stage 2 Classic is packaged in a 800g can.

Contraindications

Not suitable for those with allergies to any of the ingredients, galactosaemia or cow's milk intolerance.

Shelf life

24 months from date of manufacture.

Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

IMPORTANT NOTICE

Breastfeeding, enriched by a balanced mothers diet, is best for babies. Combined breast and bottle feeding may reduce milk supply. Deciding not to breastfeed is difficult to reverse and the social / financial implications of bottle feeding should be considered. Always follow manufacturers instructions carefully as improper use can present a health hazard or make your baby ill. Always consult your healthcare professional for advice on feeding.

Ingredients

Allergen advice see ingredients in **bold**.

Whole **Milk**, Demineralised Whey Powder (from **Milk**), Vegetable Oils (Sunflower, Coconut, Rapeseed), Lactose (from **Milk**), Galacto-oligosaccharides (from **Milk**), Skimmed **Milk** Powder, **Less than 2% of:** Whey Protein Concentrate alpha-lactalbumin (from **Milk**), Calcium Citrate, Potassium Chloride, Calcium Lactate, Sodium Citrate, Choline Bitartrate, Fructo-oligosaccharides, Magnesium Chloride, Oil from the microalgae Schizochytrium sp., Sodium Ascorbate, Potassium Hydroxide, Inositol, L-Tyrosine, Oil from Mortierella alpina, Taurine, Iron Pyrophosphate, Nucleotides (Cytidine-5'-monophosphate, disodium Uridine-5'-monophosphate, Adenosine-5'-monophosphate, disodium Inosine-5'-monophosphate, disodium Guanosine-5'-monophosphate), L-Phenylalanine, Zinc Sulphate, Vitamin E, Niacin, Calcium Pantothenate, Copper Sulphate, Thiamin, Riboflavin, Vitamin A, Vitamin B6, Manganese Sulphate, Potassium Iodide, Folic Acid, Sodium Selenite, Vitamin K, Vitamin D3, Biotin, Vitamin B12.

Typical Nutritional Values

		per 100g	per 100ml				
Energy				Thiamin	mg	0.45	0.06
	kJ	2160	279	Riboflavin	mg	1	0.13
	kcal	518	67	Niacin	mg	4.7	0.6
Fat	g	28	3.6	Vitamin B6	mg	0.37	0.05
of which:				Folate	mcg (DFE)	110	14
-saturates	g	9	1.2	Vitamin B12	mcg	0.9	0.12
-unsaturates	g	19	2.4	Pantothenic Acid	mg	3.5	0.45
Omega 3				Biotin	mcg	18	2.3
α-Linolenic Acid (ALA)	mg	390	50	Minerals			
Docosahexaenoic Acid (DHA - LCP)	mg	125	16.1	Sodium	mg	162	21
Omega 6				Potassium	mg	555	71
Linoleic Acid	mg	4185	540	Chloride	mg	362	47
Arachidonic Acid (AA – LCP)	mg	62	8	Calcium	mg	350	45
Total Carbohydrate	g	56	7.2	Phosphorous	mg	205	26
-of which sugars*	g	53	6.8	Magnesium	mg	47	6.3
-of which lactose	g	51	6.6	Iron	mg	5.8	0.75
Fibre	g	2.1	0.27	Zinc	mg	4.0	0.45
-of which Galacto - oligosaccharides	g	1.8	0.23	Copper	mg	0.41	0.053
-of which Fructo - oligosaccharides	g	0.2	0.03	Iodine	mcg	100	13
-of which 3'GL**	g	0.1	0.01	Selenium	mcg	18	2.3
Protein (N x 6.25)	g	10.3	1.3	Manganese	mcg	57	7.4
-Whey protein	g	6.0	0.77	Fluoride	mg	<0.1	<0.015
-Casein protein	g	4.0	0.52	Others			
Vitamins				Taurine	mg	45	5.8
Vitamin A	mcg-RE	400	52	Choline	mg	170	22
Vitamin D	mcg	10.5	1.35	Inositol	mg	100	13
Vitamin E	mg-α-TE	14	1.8	L- Carnitine (naturally present)	mg	6.8	0.9
Vitamin K	mcg	28	3.6	Nucleotides	mg	22	2.8
Vitamin C	mg	85	11				

*Sugars derived from milk

** 3'GL – Galactosylactose

Feeding Table

This feeding table should only be used as a guide. All babies are different and so may require more or less than is shown below. Kendamil Follow-On milk should be fed on demand.

Amount of freshly boiled and cooled water per feed	Scoops per feed
150	5
180	6
210	7

Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist.

If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.