



ESTD 1942

for healthcare professional use

Kendamil®

product data card

Organic Follow-On Milk 800g

Description

Kendamil Organic Follow On Milk is a formula uniquely made with organic whole milk fat nutrients and free from palm oil and fish oil, formulated to help meet the increased nutritional needs of infants from 6 months of age alongside a weaning diet. Exclusively manufactured in the UK using Red-Tractor and Soil Association certified organic whole milk, Kendamil is enriched with plant-based DHA and ARA, making it the only vegetarian friendly baby milk. Kendamil Follow On Milk contains a blend of GOS and HMOs in the form of 3'-GL (galactosyllactose).

Presentation

Kendamil Stage 2 Organic is packaged in a 800g can.

Contraindications

Not suitable for those with allergies to any of the ingredients, galactosaemia or cow's milk intolerance.

Shelf life

24 months from date of manufacture.

Storage

Store in a cool dry place before and after opening. Packaged in a protective atmosphere. For best before, see base of can. Use within 4 weeks of opening. Do not refrigerate and do not freeze.

IMPORTANT NOTICE

Breastfeeding, enriched by a balanced mothers diet, is best for babies. Combined breast and bottle feeding may reduce milk supply. Deciding not to breastfeed is difficult to reverse and the social / financial implications of bottle feeding should be considered. Always follow manufacturers instructions carefully as improper use can present a health hazard or make your baby ill. Always consult your healthcare professional for advice on feeding.

Ingredients

Allergen advice see ingredients in **bold**.

Organic Whole **Milk**, Organic Demineralised **Milk** Whey Protein Powder, Organic Skimmed **Milk**, Organic Vegetable Oils (Sunflower, Coconut, Rapeseed), Organic Galacto-oligosaccharides (from **Milk**), **Less than 2% of:** Calcium Citrate, Sodium Citrate, Potassium Chloride, Calcium Lactate, Magnesium Chloride, Oil from the microalgae Schizochytrium sp., Vitamin C, Potassium Hydroxide, Choline Bitartrate, Inositol, Oil from Mortierella alpina, Taurine, Iron Pyrophosphate, Nucleotides (Cytidine-5'-monophosphate, disodium Uridine-5'-monophosphate, Adenosine-5'-monophosphate, disodium Inosine-5'-monophosphate, disodium Guanosine-5'-monophosphate), Zinc Sulphate, Vitamin E, Niacin, Calcium Pantothenate, Copper Sulphate, Thiamin, Riboflavin, Vitamin A, Vitamin B6, Manganese Sulphate, Folic Acid, Potassium Iodide, Sodium Selenite, Vitamin K, Vitamin D3, Biotin, Vitamin B12.

Typical Nutritional Values

		per 100g	per 100ml				
Energy				Riboflavin	mg	1.4	0.18
	kJ	2133	275	Niacin	mg	4.7	0.6
	kcal	510	66	Vitamin B6	mg	0.4	0.05
Fat	g	27.2	3.5	Folate (DFE)	mcg (DFE)	150	19.3
of which:				Vitamin B12	mcg	1.5	0.19
-saturates	g	9.5	1.2	Pantothenic Acid	mg	3.8	0.49
-unsaturates	g	16.2	2.1	Biotin	mcg	12	1.6
Omega 3				Minerals			
α-Linolenic Acid (ALA)	mg	430	55.5	Sodium	mg	200	26
Docosahexaenoic Acid (DHA - LCP)	mg	125	16.1	Potassium	mg	580	75
Omega 6				Chloride	mg	360	46
Linoleic Acid	mg	4515	582	Calcium	mg	340	44
Arachidonic Acid (AA - LCP)	mg	62	8	Phosphorous	mg	185	23.8
Total Carbohydrate	g	54	7	Magnesium	mg	45	5.8
-of which sugars*	g	52	6.7	Iron	mg	6	0.8
-of which lactose	g	49	6.3	Zinc	mg	4.2	0.54
Fibre	g	1.8	0.23	Copper	mg	0.42	0.054
-of which Galacto - oligosaccharides	g	1.7	0.22	Iodine	mcg	100	13
-of which 3'GL**	g	0.06	0.01	Selenium	mcg	19	2.5
Protein (N x 6.25)	g	11.4	1.5	Manganese	mcg	75	9.7
Vitamins				Fluoride	mg	<0.1	<0.015
Vitamin A	mcg-RE	365	47	Others			
Vitamin D3	mcg	10.3	1.3	Taurine	mg	45	5.8
Vitamin E	mg-α-TE	10.5	1.4	Choline	mg	145	18.7
Vitamin K	mcg	28	3.6	Inositol	mg	90	11.6
Vitamin C	mg	70	9	L- Carnitine (naturally occurring)	mg	6.8	0.9
Thiamin	mg	0.5	0.06	Nucleotides	mg	21	2.7

*Sugars derived from milk

** 3'GL - Galactosyllactose

Feeding Table

This feeding table should only be used as a guide. All babies are different and so may require more or less than is shown below. Kendamil Follow-On milk should be fed on demand.

Amount of freshly boiled and cooled water per feed	Scoops per feed
150	5
180	6
210	7

Preparation

1. Wash hands, then sterilise your feeding utensils according to manufacturer's instructions.
2. Fill kettle with 1 litre of freshly run tap water (do not use repeatedly boiled water). Boil and leave to cool for 30 minutes, so it remains at a temperature of at least 70°C. Measure the required water into a sterilised bottle.
3. Using the scoop provided, add the correct number of levelled scoops to the bottle. Use the straight edge inside the lid to level each scoop.
4. Place a sterilised teat and cap on the bottle and shake well to dissolve powder.
5. Cool to a natural body temperature by running the bottle (lid on) under cold running water. Always test the temperature of the formula on the inside of your wrist.

If feed has not been used within 2 hours, we recommend that you throw it away and start again. Do not add food products such as cows' milk and do not re-heat in a microwave.